





Common Native Plants



Wild Ginger
Asarum caudatum

Distinct heart-shaped leaves form small carpets in moist shady areas.
Aromatic stems and roots.



Redwood Sorrel
Oxalis oregana

Carpets forest floor.
Flowers white to pinkish.
Leaves in 3 heart shaped sections.




Salal
Gaultheria shallon

Waxy green leaves with sticky stems.
Flowers hanging white bells.
Berries dark purple.




Thimbleberry
Rubus parviflorus

Tall shrub. Leaves 5-lobed, hairy and toothed.
Fruit red, thimble shaped.
Ripe early-mid summer.




Poison Oak
Toxicodendron diversilobum

Grows as shrub or vine.
Leaves grow in groups of three. Green in summer turning red in fall. CAUTION: Do not touch, can cause allergic reaction, skin rash.




Salmonberry
Rubus spectabilis

Common Shrub along creeks and edges of forests. Leaves serrated. Stalk has prickles.
Produces yellow to red berries.




Red Huckleberry
Vaccinium parvifolium

Deciduous, tall shrub with soft green leaves in summer.
Bark bright green.
Berries red, small, ripe in mid-summer.



Evergreen Huckleberry
Vaccinium ovatum

Tall evergreen shrub with waxy, toothed, green leaves.
Berries black, clustered, ripe in fall.




Trail Plant
Adenocaulon bicolor

Leaves smooth and green on top
Surface, fuzzy and white on underside. When trampled, the white arrow-shaped undersides point in direction of the disturber.




Common Native Trees



Tanoak

Lithocarpus densiflorus var. *densiflorus*


Leaves leathery, toothed.
Bark grayish brown.
Acorns hairy-capped and are important
Source of food for wildlife.



Red Alder

Alnus rubra


Grows in open and damp habitats.
Leaves coarsely toothed.
Bark white-gray.



California Laurel

(Pepperwood, Bay)
Umbellularia californica

Leaves shiny green, yellow in
fall, aromatic when rubbed in
hands.



Douglas-fir

Pseudotsuga menziesii var. *menziesii*


Found on ridges.
Bark brown and furrowed, brittle.
Often carries more
lichen than Coast Redwoods.



Big-leaf Maple

Acer macrophyllum


Grows along streams.
Bark covered with mosses.
Leaves yellow in fall.



Sitka Spruce

Picea sitchensis

Grows close to coast.
Needles grow on all sides of twig.
Bark gray, chip-like.



Coast Redwood

Sequoia sempervirens
(sun needles)

Fibrous reddish-brown bark, deep
furrows extend up tree. Cones
olive sized.



Coast Redwood

Sequoia sempervirens
(shade needles)

Scaly needles (left) from top of tree.
Flat needles (shown) from lower
reaches where light is
less prevalent.



Western Hemlock

(Octopus Tree)
Tsuga heterophylla

Often grows on
decaying logs and stumps.
Drooping branches with soft
needles.